

## §6.6 Logarithmic and Exponential Equations

### Properties of Exponential and Logarithmic Functions:

For  $a > 0$  and  $a \neq 1$ :

1)  $a^x = a^y$  if and only if  $x = y$ .

2) If  $x > 0$  and  $y > 0$ ,  $\log_a x = \log_a y$  if and only if  $x = y$ .

Example: Solve using the properties above.

a)  $2^x = 16$

b)  $\log_5(x+1) = \log_5(10)$

(**Note:** remember that the domain of  $y = \log_b x$  is  $(0, \infty)$ . For this reason it is always necessary to check that the solution of a logarithmic equation results in logarithms of positive numbers in the original equation.)

## Solving Exponential and Logarithmic Equations (TYPE 2)

An exponential or logarithmic equation may be solved by changing the equation into one of the following **FORMS**, where  $a$  and  $b$  are real numbers,  $a > 0$ , and  $a \neq 1$ .

- 1)  $a^{f(x)} = b$     Solve by taking the logarithms of each side. (Natural logarithms are often a good choice.)
- 2)  $\log_a f(x) = \log_a g(x)$     From the given equation,  $f(x) = g(x)$ , which is solved algebraically.
- 3)  $\log_a f(x) = b$     Solve by using the definition of logarithm to write the expression in exponential form as  $f(x) = a^b$ .

Examples: Solve the following equations for  $x$ .

a)  $4^x = 72$

d)  $\log_4 x = 3$

$$\text{b) } 4e^{2x} = 5$$

$$\text{e) } 5 + 2\ln x = 4$$